

RELATIONS BETWEEN CHILDREN AND MATERNAL NUTRIMENT IN QINGHAI AND GANSU, NORTHWESTERN RURAL CHINA

ZHANG J.,¹ KANG H.,² WANG L.F.,³ ZHAO X.Y.,⁴ LIU R.P.⁵ AND HE L.⁶

^{1,3-6}Department of Child Health Care, Xian Children's Hospital, Xian 710003, China

²Department of Pediatrics, Shaanxi Provincial People's Hospital, Xian 710068, China

ABSTRACT

Background and Objectives: We know that in low- and middle-income countries maternal and children's under nutriment is responsible for the advancement in deaths and other diseases. A study was designed here to establish a relationship between maternal and children's nutriment in northwestern rural China.

Methods: We conducted this prevalence analysis between October 2016 to December 2016 in northwestern rural China by choosing 2021 mothers of children (5 months < age ≤ 24 months) as study subjects. We used standard and calibrated methods and techniques to measure the weight and height of mothers and horizontal lengths of children.

Results: We calculated the Z-scores of 'weight for height' (WfH), 'weight for age' (WfA) and 'height for age' (HfA) and found the values less than -2 for 8.2%, 10.5% and 12.6% children, respectively. We found association between nutriment of children and maternal anthropometrical features and also between maternal characteristics and Z-scores of HfA ($P = 0.012$) and WfH ($P = 0.042$) of children was also achieved.

Conclusion: Maternal good nutriment and good salary is found to be beneficial for children's nutriment. Thus for the betterment of mother and child's health, maternal nutriment as well as her guaranteed financial empowerment should be improved.

Keywords: Maternal; children; nutriment; health; northwestern rural China.

INTRODUCTION

In recent years, it has been estimated that the under nutrition status of mother and her child is highly responsible for the one-third children disease burden and around 5 million deaths in a year. This condition is even worse in low- and middle-income countries and consequently, under nutrition status of mother and child found responsible for advancement in overall disease burden and deaths. Besides, it has also been evidenced by scientific community that maternal nutriment is very important for better health of the mother as well as for better and healthy growth and development of the child. Thus the crucial role of maternal nutriment cannot be neglected.^{1,2} In the similar fashion, association between maternal and child under nutrition with human capital and risk of adult diseases in low-income and middle-income various countries has been reviewed and revealed that damaged suffered in early life leads to premature impairment and will affect future generations.³

In general, an employed woman contributes to the household income and thus indirectly contributes to the nutrition of the family and overall, enhances her

nutriment. This happens because after getting employment a woman's power and status improve automatically and she may spend her earned money on gaining nutrition due to preferential inclusion of health and nutrition related spending from household earnings.⁴ In this way the employment status of woman boosts the child's welfare and thus helps in improving the worldwide policies of development agencies related to the development of children's health and the economic empowerment of women.⁵ Specifically, the Malaysian children's status is very bad with respect to unemployed status of mothers.⁶ Moreover, a study to reveal relationship between multiple micronutrients and nutriment among school children⁷ has helped us fight against the malnutrition problem among children.

In addition to this, the effect of maternal education on children's health has also been in explored by researchers. It was found that maternal educational status is strongly linked with the health of the children because children of less educated mothers have more malnutrition problems in comparison to those children whose mothers have good education. Hence a strong association between maternal educational status and

children's nutriment has been revealed in several countries.⁸⁻¹⁰ In addition to this, a study, related to Ghana, to establish a relationship between nutritional knowledge of mother and nutriment of the child implied that mother's knowledge about nutrition is more essential than other formal education with respect to improvement in nutriment of the child.¹¹

To the best of our knowledge, our study is the first study of its own kind to assess the relationship between maternal and children's nutriment in underdeveloped region of China *i.e.* in northwestern rural China. We aimed this work in order to fight against the malnutrition problem in this region of China.

METHODS

Survey Area and Method Design

The baseline survey area was chosen from northwestern rural China in the form of Qinghai and Gansu provinces. These provinces are mostly rural and have high land area for farming purpose. The prevalence analysis was conducted between October 2016 to December 2016. Total 2021 mothers of children (5 months < age < 24 months), living in these two provinces were chosen randomly as study subjects. We took one precaution in this selection that we chose only those mothers who were residing in the study area for more than 6 months. We got written/oral consent of all the subjects.

Questionnaire and Anthropometric Measurements

In order to estimate the socio demographic features of the mother and of the family we performed a structured pretested questionnaire for baseline survey. First we developed the questionnaire in English with the assistance of process validation and then translated it into Zhongyuan Mandarin language. We used principal component analysis for the estimation of the maternal wealth and got 22 items having 0.7 unit value of Cronbach's alpha. With the assistance of well trained staff, the height (horizontal length in case of children) and weight of both mother and children were measured during the baseline survey. We used standard and calibrated methods and techniques for anthropometric measurements. Seca medical 770 personal scale balance was used to measure the weight of barefooted children wearing light (weight) shirt (least count was 0.1 kg). With the help of a length-measuring board the horizontal length was measured to the nearest 0.1 cm.

Data Analysis

The statistical data analysis software SPSS (v16.1) was used after entering the data followed by checking the missing values and outliers. WHO Anthro software was used for the conversion of heights, ages and weights of children to respective Z-scores (*i.e.* HfA, WfH and WfA). For linear regression models variables with

respect to demography, socio-economy, education and anthropometry were selected with the help of significant Pearson's correlation coefficients. Only those variables which were significantly associated ($p < 0.200$) in the bivariate analyses were subjected to multivariate analyses in order to predict the children nutriment. In all the cases, $P < 0.050$ was considered as statistically significant.

Ethics Approval and Consent to Participate

The complete study was approved by Ethical Review Committee of Xian Children's Hospital. Besides, we also informed to the authorities in the respective localities about the purpose and objectives of our study. We also explained the advantages of our study to the mothers and then they gave oral consent. For the noble cause we referred malnourished children whose condition was severe to the health facilities in their respective localities and also informed this to the children's parents.

RESULTS

Maternal Social, Demographic, Economic and Household Characteristics

The results related to the study of maternal social, demographic and household factors with respect to rural areas of northwestern Chinese provinces: Qinghai and Gansu are shown in Table 1. All the mothers were married. Most of the mothers have Han (Muslim religion) ethnicity (72.4%) and they were not the head of the family in most of the cases (94.9%). Besides, there were other ethnicities such as Hui and Tibetan mothers were also present in effective population. More than 50% mothers were far behind their formal education (57.4%). A large number of mothers had earning based on agriculture (32.3%) but most of the mothers were unemployed (63.8%). They drink tap water (61.5%) provided by community source and 18.1% mothers had meal priority for their children. The wealth status of the families were mostly (64.3%) in medium level. A large number of families were living bamboo wall made houses (74.9%) with roof made of thatched grass (75.3%). For amusement purpose mothers listened radio in 17.9% cases and they watched television in 28.7% cases.

Anthropometric Characteristics and Relation with Nutriment

Maternal and children's anthropometric characteristic features and its relation with children's nutriment are shown in detailed Table 2. Half of the mothers (49.2%) were between the age of 26 years and 30 years, almost one-fourth (23.3%) mothers were in the range of 30-35 years and 20.4% mothers were 21-25 years old. Body mass index of most (84.7%) of the mothers were 18.6-25 kg/m² while 13.9% mothers had body mass index below 18.6 kg/m². More than half (51.8%) of the chil-

dren were female and most (65.0%) of the children were 13-24 months old. For a large number (10.2%) of children the Z-score of weight for height (WfH) were below -2. Besides, 11.7% children were below -2 Z-score of height for age (HfA) and 6.9% children were below -2 Z-score of weight for height (WfH).

Correlation between Maternal Characteristics and Children's Nutriment

Pearson correlation coefficient values with respect to establish the correlation between maternal characteristics and children's nutriment is shown in Table 3. Maternal education ($r = 0.32$, $P = 0.001$) and maternal body mass index ($r = 0.22$, $P = 0.018$) were correlated with children's Z-score of WfH. On the other hand, maternal height ($r = 0.27$, $P = 0.006$) was correlated with children's Z-score of HfA.

Anticipation of Children Nutriment

Table 4 shows the identification of anticipators of Z-scores of WfH, WfA and HfA of the children by using stepwise regression analysis. We entered all socio demographic and anthropometric features of the mothers which were significant at $P < 0.2$ into the models. The Z-score of WfH of the children was significantly predicted by normal maternal body mass index and salaried (either from private job or from government job) employment features. However, we found negative association between Z-score of WfH of the children and maternal illiteracy ($P = 0.037$). Additionally, we also found significant association between Z-score of WfA of the children and salaried employment of mother. On the other hand, a significant association was found between Z-score of HfA of the children and maternal height. We found association between Z-score of HfA of the children and tap water source in bivariate analysis but not in multivariate analysis.

DISCUSSION

We designed this baseline survey for the northwestern rural China, specifically for Qinghai and Gansu provinces. We found that normal maternal body mass index (18.6 to 25.0 kg/m²) predicted Z-score of WfH thus we can also say that children's nutriment was predicted by maternal anthropometry feature. The reason behind this outcome lies in the fact that mothers in middle and low wealth classification had lower body mass indices in comparison to mothers in high wealth classification ($P < 0.001$).

A correlation between maternal body mass index with education and income was shown in an African country's study.¹² A study to explore the micronutrient deficiencies and nutritional characteristics of diets of the children in several countries discovered that the quality of diet rather than its quantity plays the key role and hence it should be improved to overcome malnutrition.¹³ In a South-American country, Brazil signi-

ficant impact of maternal height and weight on children's nutriment was achieved when a study was performed in its day-care center.¹⁴ In our present study for northwestern rural China, we also found similar type of results in which maternal education feature was found be a positive anticipator for Z-score of WfH of the children. This can be explained on the fact that maternal education is directly proportional to their visit to healthcare centre and which is directly proportional to nutritional advice from health professionals and thus we can say that a highly educated mother is more prone to get healthy advice from professionals/doctors. Another advantage of maternal well education is that they are more feasible to get salaried employment and that will definitely enhance the purchasing and decisive power of a mother and this will directly improve children's nutriment.

In our prevalence analysis we found that Z-scores of all the three: weight for height (WfH), weight for age (WfA) and height for age (HfA) of children were strongly anticipated by salaried (either from private job or from government job) employment feature. A Brazilian cross-sectional study showed that maternal lower household income yielded lower mean Z-score of HfA among children.¹⁵ It is also important to note that besides the wealth of a household maternal owned wealth may play a crucial role as it may help her buying various food which in result improve the diet diversity of the family. Thus financial empowerment of mothers could be an essential step to improve children's nutriment. Similar result has been reported in a low income area of a Central-American country in which a correlation between Z-score of WfA and WfH of children and maternal wealth was observed⁵. However, in this study it was observed that Z-score of WfH of children was affected by their maternal employment status. Moreover, in a study in Malaysia, weight of children was positively correlated with the working hours of their mothers⁶. Besides, another study of similar kind among an African country's women and their children nutriment¹⁶ revealed that children's growth in regions was associated with working status of the mother with significant and positive impact but with high levels of malnutrition and this outcome was further complimented and supported by another study.¹⁷ Mechanism of this correlation between maternal employment and child development revealed that employed women spend significantly less time with their children's educational activities and this cannot be offset by husband.¹⁸

In our study Z-score of height for age of children was anticipated by maternal height feature (≥ 156.9 cm). This may be happened because of the fact that over the growth period children's linear growth is influenced by height of the mother as reported elsewhere.¹⁹ Another study revealed that after the adjustment of socioeconomic status, the Z-scores of HfA of

children was significantly lower when their mothers had lowest tertile of height in comparison to those children whose mothers had highest height tertile.²⁰ In our work, the Z-score of weight for height (WfH) showed that 10.2% children were underweight. Besides, Z-scores of height for age (HfA) and weight for height (WfH) showed that 11.7% children had stunting and 6.9% children had wasting. Thus we found a correlation that enhancement in maternal height was significantly associated with lower odds of stunting because we found significant enhancement in Z-score of HfA of the children with enhancement in maternal height. This result is quite consistent with a previous work on maternal nutrition studies in an African country.²¹ Interestingly, in our findings, there was a negative association between Z-score of HfA of the children and the paternal guardianship. Similar kind results were obtained previously in African countries where the mean Z-score values of females were consistently higher than those of males.²²

We found negative association between Z-scores of WfA and HfA of the children and the older maternal age (≥ 36 years age) in our study. The probable reason behind this might be the less health conscious nature of older mothers in comparison to younger mothers. It is important to note here that this is not a linear relationship as we found that youngest mothers (16 – 20) were also not health conscious. Our result is quite consistent with the World Bank report which said that children born to mid-late twenties aged mother were least prone to malnutrition.²³ Additionally, children of the teen mothers and late thirties aged mothers are significantly more likely to suffer with malnutrition.

Our prevalence analysis has large sample collection data and thus it tells the complete story of relationship between maternal and children nutriment in northwestern rural China, especially, Qinghai and Gansu provinces. Thus it would not be very difficult to generalize our results with reference to other rural provinces of northwestern China. Another advantage of our study is that it outcomes less probable causal relationships between the nutriment of mother and child.

In **conclusion**, we can say that well educated and better salaried employed mother provides a better nutriment to their children. On the other hand, Z-scores of HfA and WfH of the children were significantly associated with height of the mother and normal body mass index of the mother, respectively. The inter-connection between the nutriment of mother and child stresses that improvement in the mothers' nutriment would definitely lead to improved health outcomes of both the mothers and their children.

ACKNOWLEDGEMENTS

The authors would like to appreciate the opportunity provided by Xian Children's Hospital to conduct this research.

Conflict of Interests

The authors declare that they have no conflict of interest.

Author's Contribution

JZ: Principle author, supervised the study, HK: conceived the idea, improvement of manuscript; LFW and XYZ: data collection, literature review; RPL and LH: write-up, literature search, proof read the manuscript.

Disclaimer: None.

Financial Support: None.

REFERENCES

1. Bhutta ZA, Das JK, Rizvi A, Gaffey MF, Walker N, Horton S, et al. Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? *Lancet*, 2013; 382: 452-77.
2. Black RE, Allen LH, Bhutta ZA, Caulfield LE, Onis Md, Ezzati M, et al. Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet*, 2008; 371: 243-60.
3. Victora CG, Adair L, Fall C, Hallal PC, Martorell R, Sachdev HS, et al. Maternal and child undernutrition: consequences for adult health and human capital. *Lancet*, 2008; 371: 340-57.
4. Girma W, Genebo T. Determinants of the Nutritional Status of Mothers and Children in Ethiopia. Ethiopia Health and Nutrition Research Institute 2002. Calverton, Maryland USA. Accessed 15 Oct 2015.
5. Lamontagne JF, Engle PL, Zeitlin MF. Maternal employment, childcare, and nutritional status of 12-18-month-old children in Managua, Nicaragua, *Soc Sci Med*. 1998; 46: 403-14.
6. Shuhaimi F, Muniandy ND. The association of maternal employment status on nutritional status among children in selected kindergartens in Selangor, Malaysia. *Asian J Clin Nutr*. 2012; 4: 53-66.
7. Amare B, Moges B, Fantahun B, Tafess K, Woldeyohannes D, Yismaw G, et al. Micronutrient levels and nutritional status of school children living in Northwest Ethiopia. *Nutr J*. 2012; 11: 108-15.
8. Bbaale E. Maternal education and child nutritional status: evidence from Uganda, *Afr J Eco Mgmt Stud*. 2014; Vol. 5: pp.52-74.
9. Abubo MJ, Ndenge GK, Mwabu DK. Determinants of Children's Nutritional Status in Kenya: Evidence from Demographic and Health Surveys. *J Afr Econ*. 2008; 18: 363-87.
10. Frost MB, Forste R, Haas DW. Maternal education and child nutritional status in Bolivia: finding the links. *Soc Sci Med*. 2005; 60: 395-407.
11. Appoh LY, Krekling S. Maternal nutritional knowledge and child nutritional status in the Volta region of Ghana. *Matern Child Nutr*. 2005; 1: 100-10.
12. Fabunmi TM, Onabanjo OO, Oguntona EB, Keshinro OO, Onabanjo JA, Obanla OO, et al. Nutrient Intakes and Nutritional Status of Mothers and their Under-Five Children in a Rural Community of Oyo State, Nigeria. *Int J of Chi Hlth Nutr*. 2013; 2: 39-49.

13. Lutter CK, Rivera JA. Nutritional status of Infants and Young Children and Characteristics of Their Diets. *J Nutr.* 2003; 133: 2941S-9S.
14. Lander RL, Lander AG, Houghton L, Williams SM, Costa-Ribeiro H, Barreto DL, et al. Factors influencing growth and intestinal parasitic infections in preschoolers attending philanthropic daycare centers in Salvador, Northeast Region of Brazil. *Cad Saude Publica, Rio de Janeiro.* 2012; 28: 2177-88.
15. Felisbino-Mendes MS, Villamor E, Velasquez-Melendez G. Association of Maternal and Child Nutritional Status in Brazil: A Population Based Cross-Sectional Study. *Plos One,* 2014; 9: e87486-94.
16. Ajieroh V. A Quantitative Analysis of Determinants of Child and Maternal Malnutrition in Nigeria. Abuja, Nigeria: International Food Policy Research Institute. E-print, 2009; Accessed 3 Aug 2016.
17. Youn MJ, Leon J, Lee KJ. The influence of maternal employment on children's learning growth and the role of parental involvement. *Early Child Dev Care,* 2012; 182: 1227-46.
18. Cawley J, Liu F. Mechanisms for the Association Between Maternal Employment and Child Cognitive Development, NBER Working Papers 13609, National Bureau of Economic Research, Inc; 2007.
19. Addo OY, Stein AD, Fall CH, Gigante DP, Guntupalli AM, Horta BL, et al. Maternal height and child growth patterns. *J Pediatr.* 2013; 163: 49-554.
20. Lourenco BH, Villamor E, Augusto RA, Cardoso MA. Determinants of linear growth from infancy to school-aged years: a population-based follow-up study in urban Amazonian children. *BMC: Biomed Central Public Health,* 2012; 12: 265.
21. Gewa CA, Oguttu M, Yandell NS. Maternal nutrition in rural Kenya: health and socio-demographic determinants and its association with child nutrition. *Matern Child Nutr.* 2012; 8: 275-86.
22. Wamani H, Astrom AN, Peterson S, Tumwine JK, Tylleskar T. Boys are more stunted than girls in sub-Saharan Africa: a meta-analysis of 16 demographic and health surveys. *BMC Pediatr.* 2007; 7: 17-27.
23. World Bank: Child malnutrition. Available: <http://siteresources.worldbank.org/INTSRILANKA/ResourcesZMDG-Ch3.pdf>. Accessed 30 May 2016.